

# Literatur zum Artikel

## Molekularbiologische Grundlagen zur Pathophysiologie der Narbe

1. Aarabi S, Bhatt KA, Shi Y, et al (2007) Mechanical load initiates hypertrophic scar formation through decreased cellular apoptosis. *Faseb J* 21: 3250–3261
2. Ashcroft GS, Dodsworth J, Van Boxtel E, et al (1997) Estrogen accelerates cutaneous wound healing associated with an increase in TGF-beta1 levels. *Nat Med* 3: 1209–1215
3. Ashcroft GS, Horan MA, Ferguson MW (1998) Aging alters the inflammatory and endothelial cell adhesion molecule profiles during human cutaneous wound healing. *Lab Invest* 78: 47–58
4. Ashcroft GS, Mills SJ, Lei K, et al (2003) Estrogen modulates cutaneous wound healing by downregulating macrophage migration inhibitory factor. *J Clin Invest* 111: 1309–1318
5. Baker RH, Townley WA, Mckee S, et al (2007) Retrospective study of the association between hypertrophic burn scarring and bacterial colonization. *J Burn Care Res* 28: 152–156
6. Bayat A, McGrouther DA, Ferguson MW (2003) Skin scarring. *BMJ* 326: 88–92
7. Bloemen MC, Van Der Veer WM, Ulrich MM, et al (2009) Prevention and curative management of hypertrophic scar formation. *Burns* 35: 463–475
8. Brown DL, Kao WW, Greenhalgh DG (1997) Apoptosis down-regulates inflammation under the advancing epithelial wound edge: delayed patterns in diabetes and improvement with topical growth factors. *Surgery* 121: 372–380
9. Brown JJ, Bayat A (2009) Genetic susceptibility to raised dermal scarring. *Br J Dermatol* 161: 8–18
10. Bush JA, McGrouther DA, Young VL, et al (2011) Recommendations on clinical proof of efficacy for potential scar prevention and reduction therapies. *Wound Repair Regen* 19 (Suppl 1): s32–37
11. Butzelaar L, Ulrich MM, Berardesca E, et al (2016) Currently known risk factors for hypertrophic skin scarring: a review. *J Plast Reconstr Aesthet Surg* 69: 163–169
12. Coolen NA, Schouten KC, Middelkoop E, et al (2010) Comparison between human fetal and adult skin. *Arch Dermatol Res* 302: 47–55
13. Deitch EA, Wheelahan TM, Rose MP, et al (1983) Hypertrophic burn scars: analysis of variables. *J Trauma* 23: 895–898
14. Desmouliere A, Badid C, Bochaton-Piallat ML, et al (1997) Apoptosis during wound healing, fibrocontractive diseases and vascular wall injury. *Int J Biochem Cell Biol* 29: 19–30
15. Eming SA, Krieg T, Davidson JM (2007) Inflammation in wound repair: molecular and cellular mechanisms. *J Invest Dermatol* 127: 514–525
16. Farage MA, Miller KW, Berardesca E, et al (2009) Clinical implications of aging skin: cutaneous disorders in the elderly. *Am J Clin Dermatol* 10: 73–86
17. Farage MA, Miller KW, Elsner P, et al (2008) Intrinsic and extrinsic factors in skin ageing: a review. *Int J Cosmet Sci* 30: 87–95
18. Ferguson MW, O'kane S (2004) Scar-free healing: from embryonic mechanisms to adult therapeutic intervention. *Philos Trans R Soc Lond B Biol Sci* 359: 839–850
19. Gangemi EN, Gregori D, Berchiolla P, et al (2008) Epidemiology and risk factors for pathologic scarring after burn wounds. *Arch Facial Plast Surg* 10: 93–102
20. Gilliver SC, Ashcroft GS (2007) Sex steroids and cutaneous wound healing: the contrasting influences of estrogens and androgens. *Climacteric* 10: 276–288
21. Gilliver SC, Ruckshanthi JP, Atkinson SJ, et al (2007) Androgens influence expression of matrix proteins and proteolytic factors during cutaneous wound healing. *Lab Invest* 87: 871–881
22. Hall GK, Phillips TJ (2004) Skin and hormone therapy. *Clin Obstetr Gynecol* 47: 437–449
23. Iannello S, Milazzo P, Bordonaro F, et al (2006) Low-dose enalapril in the treatment of surgical cutaneous hypertrophic scar and keloid – two case reports and literature review. *MedGenMed* 8: 60
24. Ketchum LD, Cohen IK, Masters FW (1974) Hypertrophic scars and keloids. A collective review. *Plast Reconstr Surg* 53: 140–154
25. Kischer CW, Shetlar MR, Chvapil M (1982) Hypertrophic scars and keloids: a review and new concept concerning their origin. *Scan Electron Microsc* 1982 (Pt 4): 1699–1713
26. Knipper JA, Willenborg S, Brinckmann J, et al (2015) Interleukin-4 receptor alpha signaling in myeloid cells controls collagen fibril assembly in skin repair. *Immunity* 43: 803–816
27. Ko JH, Kim PS, Zhao Y, et al (2012) HMG-CoA reductase inhibitors (statins) reduce hypertrophic scar formation in a rabbit ear wounding model. *Plast Reconstr Surg* 129: 252e–261e
28. Kwan P, Desmouliere A, Tredget EE (2012) Molecular and cellular basis of hypertrophic scarring. In: Herndon DN (ed) *Total burn care*. Saunders Elsevier, Philadelphia, S 495–505
29. Lawrence JW, Mason ST, Schomer K, et al (2012) Epidemiology and impact of scarring after burn injury: a systematic review of the literature. *J Burn Care Res* 33: 136–146
30. Lee TJ, Jeong WS, Eom JS, et al (2013) Adjuvant chemotherapy reduces the incidence of abdominal hypertrophic scarring following immediate TRAM breast reconstruction. *Breast Cancer Res Treat* 137: 767–771
31. Longaker MT, Harrison MR, Langer JC, et al (1989) Studies in fetal wound healing: II. A fetal environment accelerates fibroblast migration in vitro. *J Pediatr Surg* 24: 793–797
32. Longaker MT, Whitby DJ, Adzick NS, et al (1990) Studies in fetal wound healing: VI. Second and early third trimester fetal wounds demonstrate rapid collagen deposition without scar formation. *J Pediatr Surg* 25: 63–68
33. Mahdavian Delavary B, Van Der Veer WM, Ferreira JA, et al (2012) Formation of hypertrophic scars: evolution and susceptibility. *J Plast Surg Hand Surg* 46: 95–101
34. Mahdavian Delavary B, Van Der Veer WM, Van Egmond M, et al (2011) Macrophages in skin injury and repair. *Immunobiology* 216: 753–762
35. Mirastschijski U, Impola U, Jahkola T, et al (2002) Ectopic localization of matrix metalloproteinase-9 in chronic cutaneous wounds. *Hum Pathol* 33: 355–364
36. Mirastschijski U, Impola U, Karsdal MA, et al (2002) Matrix metalloproteinase inhibitor BB-3103 unlike the serine proteinase inhibitor aprotinin abrogates epidermal healing of human skin wounds ex vivo. *J Invest Dermatol* 118: 55–64
37. Mirastschijski U, Jokuszies A, Vogt PM (2012) Skin wound healing: repair biology, wound and scar treatment. In: Neligan PC (ed) *Plastic surgery*. Elsevier Saunders, Philadelphia, S 267–296
38. Mirastschijski U, Sander JT, Weyand B, et al (2013) Rehabilitation of burn patients: an underestimated socio-economic burden. *Burns* 39: 262–268
39. Mirastschijski U, Sander JT, Zier U, et al (2015) The cost of post-burn scarring. *Ann Burns Fire Disasters* 28: 215–222
40. Modarressi A, Pietramaggiore G, Godbout C, et al (2010) Hypoxia impairs skin myofibroblast differentiation and function. *J Invest Dermatol* 130: 2818–2827
41. Mustoe TA, Gurtjara A (2011) The role of the epidermis and the mechanism of action of occlusive dressings in scarring. *Wound Repair Regen* 19 (Suppl 1): s16–21
42. Niessen FB, Schalkwijk J, Vos H, et al (2004) Hypertrophic scar formation is associated with an increased number of epidermal Langerhans cells. *J Pathol* 202: 121–129
43. Niessen FB, Spauwen PH, Schalkwijk J, et al (1999) On the nature of hypertrophic scars and keloids: a review. *Plast Reconstr Surg* 104: 1435–1458
44. Ogawa R (2011) Mechanobiology of scarring. *Wound Repair Regen* 19 (Suppl 1): s2–9
45. Ogawa R (2010) The most current algorithms for the treatment and prevention of hypertrophic scars and keloids. *Plast Reconstr Surg* 125: 557–568
46. Peacock EE, Jr, Madden JW, Trier WC (1970) Biologic basis for the treatment of keloids and hypertrophic scars. *South Med J* 63: 755–760

47. Quatresooz P, Hermanns JF, Paquet P, et al (2006) Mechanobiology and force transduction in scars developed in darker skin types. *Skin Res Technol* 12: 279–282
48. Rodgers KE, Ellefson DD, Espinoza T, et al (2005) Effect of NorLeu3-A(1-7) on scar formation over time after full-thickness incision injury in the rat. *Wound Repair Regen* 13: 309–317
49. Sarrazy V, Billet F, Micallef L, et al (2011) Mechanisms of pathological scarring: role of myofibroblasts and current developments. *Wound Repair Regen* 19 (Suppl 1): s10–15
50. Singer AJ, Clark RA (1999) Cutaneous wound healing. *N Engl J Med* 341: 738–746
51. Smith CJ, Smith JC, Finn MC (1987) The possible role of mast cells (allergy) in the production of keloid and hypertrophic scarring. *J Burn Care Rehabil* 8: 126–131
52. Sorg H, Tilkorn DJ, Hager S, et al (2017) Skin wound healing: an update on the current knowledge and concepts. *Eur Surg Res* 58: 81–94
53. Tomasek JJ, Gabbiani G, Hinz B, et al (2002) Myofibroblasts and mechano-regulation of connective tissue remodelling. *Nat Rev Mol Cell Biol* 3: 349–363
54. Tredget EE, Iwashina T, Scott PG, et al (1997) Determination of plasma Ntaumethylhistamine in vivo by isotope dilution using benchtop gas chromatography-mass spectrometry. *J Chromatogr B Biomed Sci Appl* 694: 1–9
55. Tredget EE, Levi B, Donelan MB (2014) Biology and principles of scar management and burn reconstruction. *Surg Clin North Am* 94: 793–815
56. Tredget EE, Shankowsky HA, Pannu R, et al (1998) Transforming growth factor-beta in thermally injured patients with hypertrophic scars: effects of interferon alpha-2b. *Plast Reconstr Surg* 102: 1317–1328
57. Van De Water L, Varney S, Tomasek JJ (2013) Mechanoregulation of the myofibroblast in wound contraction, scarring, and fibrosis: opportunities for new therapeutic intervention. *Adv Wound Care* 2: 122–141
58. Van Den Broek LJ, Niessen FB, Scheper RJ, et al (2012) Development, validation and testing of a human tissue engineered hypertrophic scar model. *ALTEX* 29: 389–402
59. Van Der Veer WM, Bloemen MC, Ulrich MM, et al (2009) Potential cellular and molecular causes of hypertrophic scar formation. *Burns* 35: 15–29
60. Van Der Wal MB, Vloemans JF, Tuinebreijer WE, et al (2012) Outcome after burns: an observational study on burn scar maturation and predictors for severe scarring. *Wound Repair Regen* 20: 676–687
61. Walraven M, Akershoek JJ, Beelen RH, et al (2017) In vitro cultured fetal fibroblasts have myofibroblast-associated characteristics and produce a fibrotic-like environment upon stimulation with TGF-beta1: is there a thin line between fetal scarless healing and fibrosis? *Arch Dermatol Res* 309: 111–121
62. Walraven M, Gouverneur M, Middelkoop E, et al (2014) Altered TGF-beta signaling in fetal fibroblasts: what is known about the underlying mechanisms? *Wound Repair Regen* 22: 3–13
63. Walraven M, Talhout W, Beelen RH, et al (2016) Healthy human second-trimester fetal skin is deficient in leukocytes and associated homing chemokines. *Wound Repair Regen* 24: 533–541
64. Wang J, Dodd C, Shankowsky HA, et al (2008) Deep dermal fibroblasts contribute to hypertrophic scarring. *Lab Invest* 88: 1278–1290
65. Wang J, Jiao H, Stewart TL, et al (2007) Improvement in postburn hypertrophic scar after treatment with IFN-alpha2b is associated with decreased fibrocytes. *J Interferon Cytokine Res* 27: 921–930
66. Willenborg S, Eckes B, Brinckmann J, et al (2014) Genetic ablation of mast cells redefines the role of mast cells in skin wound healing and bleomycin-induced fibrosis. *J Invest Dermatol* 134: 2005–2015
67. Willenborg S, Eming SA (2014) Macrophages – sensors and effectors coordinating skin damage and repair. *JDDG* 12: 214–221, 223
68. Wong VW, Akaishi S, Longaker MT, et al (2011) Pushing back: wound mechanotransduction in repair and regeneration. *J Invest Dermatol* 131: 2186–2196
69. Wray RC (1983) Force required for wound closure and scar appearance. *Plast Reconstr Surg* 72: 380–382
70. Yang L, Scott PG, Giuffre J, et al (2002) Peripheral blood fibrocytes from burn patients: identification and quantification of fibrocytes in adherent cells cultured from peripheral blood mononuclear cells. *Lab Invest* 82: 1183–1192
71. Zhu Z, Ding J, Ma Z, et al (2016) Systemic depletion of macrophages in the subacute phase of wound healing reduces hypertrophic scar formation. *Wound Repair Regen* 24: 644–656
72. <https://www.verbrennungsmedizin.de/verbrennungsregister/verbrennungsregister-projekt.php>