

Die Gefäßdarstellung bei Perforator-Lappenplastiken

Welche Technik ist sinnvoll?

CORNELIUS DIETER SCHUBERT¹, TOBIAS HIRSCH¹, MARCUS LEHNHARDT¹, ADRIEN DAIGELE²

BOCHUM, TÜBINGEN

1. Stevenson TR, Rubin JM, Herzenberg JE (1988) Vascular patency of fibular free graft: assessment by Doppler color-flow imager: a case report. *J Reconstr Microsurg* 4: 409–413
2. Chang BW, Luethke R, Berg WA, et al (1994) Two-dimensional color Doppler imaging for precision preoperative mapping and size determination of TRAM flap perforators. *Plast Reconstr Surg* 93: 197–200
3. Chidyllo SA, Jacobs JS (1993) The application of duplex ultrasonography in the preoperative evaluation of patients prior to TRAM flap reconstruction. *Plast Reconstr Surg* 92: 174–175
4. Rand RP, Cramer MM, Strandness DE Jr (1994) Color-flow duplex scanning in the preoperative assessment of TRAM flap perforators: a report of 32 consecutive patients. *Plast Reconstr Surg* 93: 453–459
5. Hallock GG (1994) Evaluation of fasciocutaneous perforators using color duplex imaging. *Plast Reconstr Surg* 94: 644–651
6. Hallock GG, Brown CL, DeSantis MJ (1995) Preoperative identification of perforating vessels to the skin by color duplex imaging. *J Ultrasound Med* 14: 263–266
7. Schwabegger AH, Bodner G, Rieger M, et al (1999) Internal mammary vessels as a model for power Doppler imaging of recipient vessels in microsurgery. *Plast Reconstr Surg* 104: 1656–1665
8. Loh N, Ch'en IY, Olcott E, et al (1997) Power Doppler imaging in preoperative planning and postoperative monitoring of muscle flaps. *J Clin Ultrasound* 25: 465–471
9. Anderson T, McDicken WN (2002) The difference between Colour Doppler Velocity imaging and Power Doppler imaging. *Eur J Echocardiogr* 3: 240–244
10. Knobloch K, Reuss E, Gohritz A, et al (2009) Umfrage zu Methoden der präoperativen Lokalisation von Perforans-Gefäßen zur erleichterten Planung von Perforans-Lappenplastiken. *Handchir Mikrochir Plast Chir* 41: 322–326
11. Smit JM, Klein S, Werker PM (2010) An overview of methods for vascular mapping in the planning of free flaps. *J Plast Reconstr Aesthet Surg* 63: e674–e682
12. Pratt GF, Rozen WM, Chubb D, et al (2012) Preoperative imaging for perforator flaps in reconstructive surgery: a systematic review of the evidence for current techniques. *Ann Plast Surg* 69: 3–9
13. Saber AA, Meslemi AM, Davis R, Pimentel R (2004) Safety zones for anterior abdominal wall entry during laparoscopy: a CT scan mapping of epigastric vessels. *Ann Surg* 239: 182–185
14. Klasson S, Svensson H, Malm K, et al (2015) Preoperative CT angiography versus Doppler ultrasound mapping of abdominal perforator in DIEP breast reconstructions: a randomized prospective study. *J Plast Reconstr Aesthet Surg* 68: 782–786
15. Rozen WM, Garcia-Tutor E, Alonso-Burgos A, et al (2010) Planning and optimising DIEP flaps with virtual surgery: the Navarra experience. *J Plast Reconstr Aesthet Surg* 63: 289–297
16. Masia J, Clavero JA, Larranaga JR, et al (2006) Multidetector-row computed tomography in the planning of abdominal perforator flaps. *J Plast Reconstr Aesthet Surg* 59: 594–599
17. Masia J, Clavero JA, Larranaga JR, et al (2008) Preoperative planning of the abdominal perforator flap with multidetector row computed tomography: 3 years of experience. *Plast Reconstr Surg* 122: 80e–81e
18. Masia J, Larranaga J, Clavero JA, et al (2008) The value of the multidetector row computed tomography for the preoperative planning of deep inferior epigastric artery perforator flap: our experience in 162 cases. *Ann Plast Surg* 60: 29–36
19. Clavero JA, Masia J, Larranaga J, et al (2008) MDCT in the preoperative planning of abdominal perforator surgery for post-mastectomy breast reconstruction. *AJR* 191: 670–676
20. Hamdi M, Van Landuyt K, Van Hedent E, Duyck P (2007) Advances in autogenous breast reconstruction: the role of preoperative perforator mapping. *Ann Plast Surg* 58: 18–26
21. Kosutic D, Pejkovic B, Anderhuber F, et al (2012) Complete mapping of lateral and medial sural artery perforators: anatomical study with Duplex-Doppler ultrasound correlation. *J Plast Reconstr Aesthet Surg* 65: 1530–1536
22. Brenner DJ, Hall EJ (2007) Computed tomography – an increasing source of radiation exposure. *N Engl J Med* 357: 2277–2284
23. Taylor GI, Doyle M, McCarten G (1990) The Doppler probe for planning flaps: anatomical study and clinical applications. *Br J Plast Surg* 43: 1–16
24. Figus A, Wade RG, Gorton L, et al (2012) Venous perforators in DIEAP flaps: an observational anatomical study using duplex ultrasonography. *J Plast Reconstr Aesthet Surg* 65: 1051–1059

25. Gravvanis A, Tsoutsos D, Papanikolaou G, et al (2014) Refining perforator selection for deep inferior epigastric perforator flap: the impact of the dominant venous perforator. *Microsurgery* 34: 169–176
26. Rozen WM, Phillips TJ, Ashton MW, et al (2008) Preoperative imaging for DIEA perforator flaps: a comparative study of computed tomographic angiography and doppler ultrasound. *Plast Reconstr Surg* 121: 1–8
27. Scott JR, Liu D, Said H, et al (2010) Computed tomographic angiography in planning abdomen-based microsurgical breast reconstruction: a comparison with color duplex ultrasound. *Plast Reconstr Surg* 125: 446–453
28. Blondeel PN, Beyens G, Verhaeghe R, et al (1998) Doppler flowmetry in the planning of perforator flaps. *Br J Plast Surg* 51: 202–209
29. Seidenstucker K, Munder B, Richrath P, et al (2010) A prospective study using color flow duplex ultrasonography for abdominal perforator mapping in microvascular breast reconstruction. *Med Sci Monit* 16: MT65–70
30. Yu P, Youssef A (2006) Efficacy of the handheld Doppler in preoperative identification of the cutaneous perforators in the anterolateral thigh flap. *Plast Reconstr Surg* 118: 928–933
31. Stekelenburg CM, Sonneveld PM, Bouman MB, et al (2014) The hand held Doppler device for the detection of perforators in reconstructive surgery: what you hear is not always what you get. *Burns* 40: 1702–1706
32. Dorfman D, Pu LL (2014) The value of color duplex imaging for planning and performing a free anterolateral thigh perforator flap. *Ann Plast Surg* 72 (Suppl 1): S6–8
33. Iida H, Ohashi I, Kishimoto S, et al (2003) Preoperative assessment of anterolateral thigh flap cutaneous perforators by colour Doppler flowmetry. *Br J Plast Surg* 56: 21–25
34. Tsukino A, Kurachi K, Inamiya T, Tanigaki T (2004) Preoperative color Doppler assessment in planning of anterolateral thigh flaps. *Plast Reconstr Surg* 113: 241–246
35. Cheng HT, Lin FY, Chang SC (2013) Diagnostic efficacy of color Doppler ultrasonography in preoperative assessment of anterolateral thigh flap cutaneous perforators: an evidence-based review. *Plast Reconstr Surg* 131: 471e–473e
36. Ensat F, Babl M, Conz C, et al (2012) The efficacy of color duplex sonography in preoperative assessment of anterolateral thigh flap. *Microsurgery* 32: 605–610
37. Khan UD, Miller JG (2007) Reliability of handheld Doppler in planning local perforator-based flaps for extremities. *Aesthetic Plast Surg* 31: 521–525
38. Jakubietz RG, Jakubietz MG, Grünert JG, et al (2011) Propeller-Lappenplastiken: die Reliabilität der präoperativen, unidirektionale Dopplersonografie. *Handchir Mikrochir Plast Chir* 43: 76–80.
39. Daigeler A, Schubert C, Hirsch T, Behr B, Lehnhardt M (2018) Farbkodierte Duplexsonographie und „Power-Doppler. Erhöhung der Patientensicherheit durch effiziente Planung. *Handchir Mikrochir Plast Chir* (im Druck)